

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

---

2014

# Greenhouse Gas Emissions from Soil Treated with Dung Pats

Kenneth Evans

*University of Nebraska-Lincoln*

Ana Wingeyer

*Instituto Nacional de Tecnología Agropecuaria*

Martha Mamo

*University of Nebraska-Lincoln, mmamo3@unl.edu*

Walter H. Schacht

*University of Nebraska-Lincoln, wschacht1@unl.edu*

Pamela Sutton

*University of Nebraska-Lincoln*

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Botany Commons](#), [Horticulture Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

---

Evans, Kenneth; Wingeyer, Ana; Mamo, Martha; Schacht, Walter H.; Sutton, Pamela; and Eskridge, Kent M., "Greenhouse Gas Emissions from Soil Treated with Dung Pats" (2014). *Agronomy & Horticulture -- Faculty Publications*. 756.

<http://digitalcommons.unl.edu/agronomyfacpub/756>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Kenneth Evans, Ana Wingeyer, Martha Mamo, Walter H. Schacht, Pamela Sutton, and Kent M. Eskridge

## Rationale

Change in grazing density can influence dung distribution patterns, with potential impacts on the abundance and frequency of dung beetles populations and nutrient cycling in grazing systems. The goal of this research was to quantify and characterize the fate of nutrient pulses from cow dung after deposition, and the associated effects of dung beetle activity. Mass and nutrient loss of dung, changes in soil nutrients below and around dung pats, and fluxes of greenhouse gases (GHGs) were monitored overtime. The results on GHG fluxes are presented here and another paper by Evans et al. presents soil results.

## Hypotheses

Dung beetles activity can affect the amount and timing of pulses of GHGs emitted from decomposing dung pats.

- Dung pats exposed to dung beetle activity will emit higher fluxes of CO<sub>2</sub> and N<sub>2</sub>O, and lower fluxes of CH<sub>4</sub> than those not exposed to dung beetles.
- Dung pats exposed to dung beetle activity will exhibit peak fluxes of CO<sub>2</sub> earlier and peak fluxes of N<sub>2</sub>O later than those not exposed to dung beetles.

## Materials & Methods

### Site Description

Research was conducted at the University of Nebraska-Lincoln, Barta Brothers Ranch (42° 13'28.65"N, 99° 38'19.17"W) on subirrigated, sandy to fine sandy loam soils in the Valentine series.

### Experimental Design and Treatments

- Three treatments were arranged in a repeated measurement RCB with 8 blocks and replicated during grazing season (sequential June and July experiments).
- Treatments included artificially created 20 cm diameter pats from 1.5 L of homogenized beef cattle manure placed directly on the ground (BEETLE), inside a wire-mesh exclusion cage (NO BEETLE), and a no dung treatment (CONTROL).
- Gas samples were taken at 1, 2, 3, 7, 10, 14, 21, 28, and 56 days after dung pat application (DAA).



### Measurements and Analyses


- GHGs sampling followed GraceNet protocols for chamber method (Parkin and Venterea, 2010), with collection in 10 min. intervals for up to 30 min. on specified DAA.




- Samples were analyzed for CO<sub>2</sub>-C, N<sub>2</sub>O-N, and CH<sub>4</sub>-C concentrations using a Varian GC-450.
- Soil temperature and moisture at 10 and 20 cm depths, air temperature, and precipitation were monitored continuously a weather station.
- Fluxes of GHGs were calculated from regression analysis for each DAA (Parker and Venterea, 2010).
- Generalized linear mixed model with repeated measures. Separation of treatments were done using LSM (alpha=0.05).

## Results

### Weather and Soil Water Content - June experiment (28 DAA)



### N<sub>2</sub>O-N Flux




### June

- Sampling time was significant.

### July

- Treatment by sampling time interaction was significant
  - 10 DAA: NO BEETLE = control; NO BEETLE > BEETLE; BEETLE < control
  - 28 DAA: NO BEETLE = control; NO BEETLE < BEETLE; BEETLE = control
  - 1, 2, 3, 7, 14, 21, and 56 DAA: NO BEETLE = BEETLE = control

### CH<sub>4</sub>-C Flux



### June

- Sampling time was significant

### July

- No significant effects.

### Dung Beetle Colonization of dung pats

Treatment	Dung Beetles
NO BEETLE	0.5 ± 0.5 b
BEETLE	8.5 ± 1.5 a

Mean ± se dung beetle number per dung pat. n=8.


At 1, 2, and 3 DAA dung beetle counts were significantly less in the NO BEETLE than in the BEETLE treatment.

Placing dung pats inside wire mesh cages effectively excluded dung beetles.

Dung beetle abundance was tested using flotation plus manual search on dung pats harvested at 1, 2, and 3 DAA



### CO<sub>2</sub>-C Flux



### June (wetter)

- Treatment and timing by sampling time interactions were significant.
  - 1 DAA: NO BEETLE = BEETLE; NO BEETLE > control; BEETLE > control
  - 3 DAA: NO BEETLE = BEETLE; NO BEETLE > control; BEETLE > control
  - 7 DAA: NO BEETLE = BEETLE; NO BEETLE = control; BEETLE > control
  - 10 and 14 DAA: NO BEETLE = BEETLE; NO BEETLE > control; BEETLE = control
  - 2, 21, 28, and 56 DAA: NO BEETLE = BEETLE = control

### July

- Treatment and interactions were not significant



June experiment field layout (left), dung beetles on a BEETLE dung pat (top).

## Summary and Conclusions

- Treatment was significant for CO<sub>2</sub>-C flux but not for N<sub>2</sub>O-N and CH<sub>4</sub>-C fluxes
- CO<sub>2</sub>-C flux was highest in NO BEETLE TRT 5 out of the 9 sampling dates.
- There was no consistent trend in GHG flux when pat was covered to exclude dung beetles and flux was similar between NO BEETLE and BEETLE TRTs.
- Contrary to Pentilla et al., 2013 and initial hypothesis, June CO<sub>2</sub>-C flux was most often highest from NO BEETLE treatment and there was no significant difference in treatments during July.
- In accordance with Pentilla et al., 2013 and initial hypothesis, July peak N<sub>2</sub>O-N flux occurred at later DAA in BEETLE compared to control and NO BEETLE.

## Acknowledgement

This project was supported by Agriculture and Food Research Initiative Competitive Grant or National Integrated Water Quality Grant Program no. 2013-67019-21394 from the USDA National Institute of Food and Agriculture. We would like to thank Jon Soper, Heidi Hillhouse, Pat Wagner, Matt Judkins, Erin Hatch, Torie Lindsey, and Jenna Beckman for their assistance in field and laboratory work.

## References

- Parkin, T., and R. Venterea. 2010. Chamber-Based Trace Gas Flux Measurements, chapter 3. USDA-ARS GRACEnet Protocols.
- Pentilla, Atte, E.M. Slade, A. Simojoki, T. Riutta, K. Minkinen, T. Roslin (2013) Quantifying Beetle-Mediated Effects on Gas Fluxes From Dung Pats. Plos One, Vol. 8 (8), pp. 1-7.